

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 31/69, 38/55</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/56753</b> <b>(43) International Publication Date:</b> 11 November 1999 (11.11.99)
<b>(21) International Application Number:</b> PCT/US99/09652 <b>(22) International Filing Date:</b> 3 May 1999 (03.05.99) <b>(30) Priority Data:</b> 60/084,128 4 May 1998 (04.05.98) US <b>(71) Applicant:</b> POINT THERAPEUTICS, INC. [US/US]; 75 Kneeland Street, Boston, MA 02111 (US). <b>(72) Inventors:</b> WALLNER, Barbara, P.; 64 Arrowhead Road, Weston, MA 02193 (US). JONES, Barry; 80 Wendell Street, Cambridge, MA 02138 (US). MILLER, Glenn, T.; 112 Montclair Road, Haverhill, MA 01830 (US). ADAMS, Sharlene; 170 Sycamore Street, Watertown, MA 02172 (US). <b>(74) Agent:</b> PLUMER, Elizabeth, R.; Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> HEMATOPOIETIC STIMULATION  <b>(57) Abstract</b>  Methods and products for stimulating hematopoietic, preventing low levels of hematopoietic cells and producing increased numbers of hematopoietic and, mature blood cells are provided. The methods and products can be used both <i>in vivo</i> and <i>in vitro</i> .		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

- 1 -

## **HEMATOPOIETIC STIMULATION**

### **Background of the Invention**

The present invention relates to methods and products for producing increased  
5 numbers of hematopoietic cells, of restoring to preselected normal levels numbers of  
hematopoietic cells, to therapies for treating deficiencies in hematopoietic cells and to *in vitro*  
methodologies for culturing hematopoietic cells.

PT-100 is a dipeptide consisting of valine-prolineboronic acid (ValboroPro) designed  
to interact with the cell surface receptor CD26. CD26, a type II transmembrane protein is  
10 expressed on the cell surface of a number of cell types, including lymphocytes (Marguet, D. et  
al., *Advances in Neuroimmunol.* 3:209-215 (1993)), hematopoietic cells (Vivier, I. et al., *J.*  
*Immunol.* 147:447-454 (1991); Bristol, et al., *J. Immunol.* 149:367 (1992)) thymocytes (Dang,  
N.H. et al., *J. Immunol.* 147:2825-2832 (1991), Tanaka, T. et al., *J. Immunol.* 149:481-486  
(1992), Darmoul, D. et al., *J. Biol. Chem.* 267:4824-4833 (1992)), intestinal brush border  
15 membrane and endothelial cells. Cell surface associated CD26 is a sialoglycoprotein, with  
most of its mass on the outside of the cell.

CD26 has been best characterized on peripheral T cells where it functions as a potent  
costimulatory signal for T cell activation. Its surface expression is upregulated upon T cell  
activation (Dong, R.P. et al., *Cell* 9:153-162 (1996), Torimoto, Y. et al., *J. Immunol.* 147:2514  
20 (1991), Mittrucker, H-W. et al., *Eur. J. Immun.* 25:295-297 (1995), Hafler, D.A. et al, *J.*  
*Immunol.* 142:2590-2596 (1989), Dang, N.H. et al., *J. Immunol.* 144:409 (1990)). CD26 has  
also been identified in rodents as an important regulatory surface receptor in hematopoiesis  
and lymphoid development (Vivier, I. et al., *J. Immunol.* 147:447-454 (1991)). The primary  
structure of CD26 is highly conserved between species (Ogata, S. et al., *J. Biol. Chem.*  
25 264:3596-3601 (1998)). In humans CD26 seems to be involved in the regulation of  
thymocyte activation, differentiation and maturation (Dang, N.H. et al., *J. Immunol.*  
147:2825-2832 (1991); Kameoka, J. et al., *Blood* 85:1132-1137 (1995)). We have evidence  
that CD26 is expressed within the human and murine hematopoietic systems.

CD26 is an ectoenzyme with activity identical to that of Dipeptidyl Peptidase IV  
30 (DPP-IV), a serine type exopeptidase with high substrate specificity. It cleaves N-terminal  
dipeptides from proteins if the penultimate amino acid is proline. or in some cases alanine  
(Fleischer, B. *Immunol. Today* 15:180 (1994)). PT-100 is a potent inhibitor of DPP-IV

- 2 -

activity.

The prior art PCT published application WO94/03055 teaches methods of producing increased numbers of hematopoietic cells by administering inhibitors of DPP-IV. The teaching of this published application, however, is that dosages of at least 1mg/kg body weight are necessary to achieve such hematopoietic cell increases. This published application also teaches that inhibitors are administered to mammals which have an established deficiency of hematopoietic cells. The teaching also suggests that cytokines be administered in conjunction with the inhibitors to increase the production of hematopoietic cells in a subject.

#### **Summary of the Invention**


The invention is based upon a variety of surprising and unexpected findings. It has been discovered, unexpectedly, that the agents useful according to the invention stimulate growth factor production by stromal cells. It also has been discovered, unexpectedly, that the agents useful according to the invention stimulate proliferation of primitive hematopoietic progenitor cells, but do not stimulate directly the differentiation or proliferation of committed progenitor cells. It further has been discovered, unexpectedly, that the agents useful according to the invention can be administered at doses much lower than would have been expected according to the teachings of the prior art. Another unexpected finding is that the agents according to the invention can accelerate the time it takes to achieve hematopoietic cell recovery after treatment with an hematopoietic cell inhibitor. Another unexpected finding is that the agents useful according to the invention can at relatively low doses, restore normal levels of neutrophils at least as fast as the most successful commercially available product used worldwide for this purpose, except that the agents useful according to the invention can be used orally, whereas the commercially available product (which represents more than a billion dollar market) must be injected. These unexpected results have important therapeutic and experimental research implications.

According to one aspect of the invention, a method is provided for treating a subject to stimulate hematopoiesis in the subject. The invention involves administering to a subject in need of such treatment an amount of an agent effective to increase the number of hematopoietic cells or mature blood cells in the subject, wherein the amount is less than 1 mg/kg body weight per day and wherein the agent is a compound of Formula I.

The agents useful according to the invention are compounds of Formula I:


- 3 -

Formula I



wherein m is an integer between 0 and 10, inclusive; A and A<sub>1</sub> are L-amino acid residues (for glycine there is no such distinction) such that the A in each repeating bracketed unit can be a different amino acid residue; the C bonded to B is in the L-configuration; the bonds between A and N, A<sub>1</sub> and C, and between A<sub>1</sub> and N are peptide bonds; and each X<sub>1</sub> and X<sub>2</sub> is, independently, a hydroxyl group or a group capable of being hydrolysed to a hydroxyl group in aqueous solution at physiological pH. By "the C bonded to B is in the L-configuration" is meant that the absolute configuration of the C is like that of an L-amino acid.

Thus, the



group has the same relationship to the C as the --COOH group of an L-amino acid has to its α carbon. In some embodiments, A and A<sub>1</sub> are independently proline or alanine residues; m is 0; X<sub>1</sub> and X<sub>2</sub> are hydroxyl groups; the inhibitor is L-Ala-L-boroPro; and the inhibitor is L-Pro-L-boroPro.

In one important aspect of the invention, the subject has an abnormally low level of hematopoietic cells or mature blood cells and the agent is administered in an amount effective to restore levels of a hematopoietic cell-type or mature blood cell-type to a preselected normal or protective level. The agent preferably is administered to the subject in at least 2 doses in an 18 hours period. The invention has particularly important applications in the restoration of normal or protective levels of neutrophils, erythrocytes and platelets. The most preferred agent is ValBoroPro.

According to another aspect of the invention, a method is provided for shortening or eliminating the time that a subject has an abnormally low level of hematopoietic or mature blood cells resulting from treatment with a hematopoietic cell inhibitor. An agent is administered to a

- 4 -

subject in need of such treatment in an amount effective to increase the number of hematopoietic cells or mature blood cells in the subject, wherein the administration of the agent begins prior to or substantially simultaneous with administration of the hematopoietic cell inhibitor. The agents and the preferred agent are as described above. In one important embodiment, the hematopoietic cell inhibitor causes an abnormally low level of hematopoietic cells or mature blood cells in the subject and the agent is administered in an amount effective to restore levels of a hematopoietic cell type to a preselected normal or protective level. Preferably, the agent is administered to the subject in at least 2 doses in an 18 hour period. In important embodiments, the agent is used to restore in the subject normal or protective levels of neutrophils, erythrocytes or platelets. The preferred effective amount of agent is as described above.

According to another aspect of the invention, a method is provided for preparing a subject for treatment with a hematopoietic cell inhibitor. The method involves administering to the subject prior to the subject receiving the hematopoietic cell inhibitor an agent in an amount effective to stimulate in the subject production of growth factors. In one embodiment the agent stimulates stromal cell production of growth factor. The agents and the preferred agent are as described above. In one important embodiment, the growth factor is granulocyte colony stimulating factor. In other embodiments the growth factor is selected from the group consisting of IL-1, IL-2, IL-3, IL-4, IL-6, IL-11, IL-17, TPO, EPO, MCSF, GMCSF, FLT-3 Ligand and Stem Cell Factor. Preferably, the amount administered to the subject is less than 1 mg/kg body weight per day. It also is preferred that the administration of the agent be in at least 2 doses of the agent in an 18 hour period.

According to another aspect of the invention, a method is provided for treating a subject to increase the number of hematopoietic cells or mature blood cells in the subject. An agent is administered to a subject in need of such treatment in an amount effective to increase hematopoietic cells or mature blood cells in the subject, wherein the agent is administered in a first regimen consisting of 2 doses or 3 doses in an 18 hour period. The agents and the preferred agent are as described above. In one important embodiment, the agent is administered in a second regimen consisting of 2 doses or 3 doses in an 18 hours period, wherein the second regimen is separate in time from the first regimen. In another embodiment, the agent is administered in a third regimen consisting of 2 doses or 3 doses in an 18 hour period, wherein the third regimen is separate in time from the first and second regimens. In other embodiments, the agent is administered optionally in a fourth regimen, a fifth regimen, a sixth regimen, or a seventh

- 5 -

regimen, wherein each of such regimens consists of 2 doses or 3 doses in an 18 hours period, and wherein the regimens are separate in time from one another and from the prior regimens. In one important embodiment, the subject has an abnormally low neutrophil count and the amount is effective to restore in the subject a preselected level of neutrophils. In other important  
5   embodiments the subject has abnormally low levels of erythrocytes and platelets. The preferred dosages, agents, and the like are as described above. In important embodiments, the dosage is no more than six regimens, no more than five regimens, no more than four regimens, no more than three regimens, and even no more than two regimens.

According to another aspect of the invention, a method is provided for preparing a  
10   subject's cells for reintroduction into the subject. The method involves treating the subject with an agent in an amount effective to stimulate in the subject the hematopoietic cells, then collecting the hematopoietic cells from the subject. The collected cells later are reintroduced into the subject. The collected cells optionally can be *ex vivo* cultured. The agents and preferred agent are as described above. In one embodiment, the *ex vivo* culturing is carried out in the presence  
15   of an amount of the agent effective to stimulate proliferation of the collected cells. In another embodiment, the concentration of the agent in medium surrounding the collected cells is less than  $10^{-8}$  moles per liter, and less than  $10^{-9}$  moles per liter and even less than  $10^{-10}$  moles per liter.

According to another aspect of the invention, a method is provided for stimulating growth factor production by stromal cells. The method involves contacting the stromal cells with an  
20   agent in an amount effective to stimulate growth factor production by the stromal cells. The agents and the preferred agent are as described above. In one embodiment, the stromal cells are in an *in vitro* layer of stromal cells for supporting early progenitor cell growth and further comprising culturing the stem cells in the presence of these stromal cells. In another embodiment, the stromal cells are *in vivo* in a subject. In another embodiment, the growth factor  
25   is granulocyte colony stimulating factor. In other embodiments the growth factor is selected from the group consisting of IL-1, IL-2, IL-3, IL-4, IL-6, IL-11, IL-17, TPO, EPO, MCSF, GMCSF, FLT-3 Ligand and Stem Cell Factor. In an *in vivo* embodiment, the agent is administered to a subject in an amount less than 1 mg/kg body weight per day. In still another embodiment, the stem cells are cultured in an environment free of exogenously added granulocyte colony  
30   stimulating factor. In important embodiments the stromal cells are bone marrow or thymic stromal cells.

According to another aspect of the invention, a kit is provided for treating a subject having

- 6 -

an abnormally low level of hematopoietic cells resulting from treatment with a hematopoietic cell inhibitor or for treating prophylactically a subject being treated with a hematopoietic cell inhibitor to prevent decrease or loss of hematopoietic and/or mature blood cells. The kit is a package containing a first dosage and instructions for treating a subject substantially simultaneous with  
5 or prior to treatment with the hematopoietic cell inhibitor. The package also contains a second dosage and instructions for treating a subject only after treatment with the hematopoietic cell inhibitor. The dosages are in effective amounts and the agents and preferred agent are as described above. In one embodiment, the second dosage is between 2 and 5 regimens, each of the regimens consisting of 2 or 3 doses per day of the agent. In one embodiment, the combination  
10 of the doses is less than 1 mg/kg body weight per day. One preferred kit is for treatment of neutropenia. Other preferred kits are for treatment of an abnormally low level of erythrocytes or platelets.

According to still another aspect of the invention, a kit is provided for treating a subject having abnormally low level of hematopoietic cells. The kit is a package containing a complete  
15 dosage for restoring normal levels of a hematopoietic cell type. The package consists essentially of: (1) a first dosage in an effective amount for administration to the subject during a first day, (2) a second dosage in an effective amount for administration to the subject during a second day, (3) optionally, a third dosage in an effective amount for administration to the subject during a third day, (4) optionally, a fourth dosage in an effective amount for administration to the subject during  
20 a fourth day, (5) optionally, a fifth dosage in an effective amount for administration to the subject during a fifth day, (6) optionally, a sixth dosage in an effective amount for administration to the subject during a sixth day and (7) optionally, a seventh dosage in an effective amount for administration to the subject during a seventh day. The agents and preferred agent are as described above. In one important embodiment, each of the dosages consists of 2 or 3 doses of  
25 the agent for administration each day. Preferred doses and dosages are as described above. In important embodiments, the kit consists essentially of less than 5, less than 4, and less than 3 and even less than 2 dosages.

These and other aspects of the invention will be described in greater detail below.

#### **Brief Description of the Drawings**

30 Figure 1 is a medicinal package for administering a 5 day medicinal course of treatment for treating myelosuppression or anemia resulting from cancer chemotherapy.

Figure 2 is a graph depicting the regeneration of neutrophils in cyclophosphamide-treated



- 7 -

mice. PT-100 at indicated doses or saline administered by gavage. Absolute neutrophil counts in mice not treated with cyclophosphamide are on the average  $190 \times 10^4$  cells/ml as indicated by the dashed horizontal line.

Figure 3 is a graph depicting the regeneration of neutrophils in cyclophosphamide-treated mice in response to subcutaneous administration of PT-100. Saline or PT-100 were administered b.i.d. for 5 consecutive days. The average absolute neutrophil count in mice not treated with cyclophosphamide was  $185 \times 10^4$  cells/ml as indicated by the horizontal dashed line.

Figure 4 is a graph depicting the regeneration of neutrophils in cyclophosphamide-treated mice in response to PT-100 and granulocyte colony stimulating factor. PT-100 in saline was administered by gavage, and GCSF by subcutaneous injections, for 5 days. Absolute neutrophil count in mice not treated with cyclophosphamide are on average  $190 \times 10^4$  cells/ml as indicated by the dashed horizontal line.

Figure 5 is a graph depicting the effect of PT-100 dose number on the regeneration of neutrophils in cyclophosphamide-treated mice. PT-100, at indicated concentrations, was administered either once or twice per day subcutaneously for 5 days. The average absolute neutrophil count for mice not treated with cyclophosphamide was  $200 \times 10^4$  cells/ml as indicated by the dashed horizontal line.

Figure 6 is a graph depicting the effect of duration of PT-100 administration on absolute neutrophil count and rate of neutrophil recovery in cyclophosphamide-treated mice. PT-100 ( $5 \mu\text{g/b.i.d.}$ ) was administered to cyclophosphamide-treated mice by gavage for the indicated length of time. The dashed horizontal line indicates the average absolute neutrophil count for mice not treated with cyclophosphamide.

Figure 7 is a graph showing the effect of duration of PT-100 treatment on the regeneration of neutrophils in cyclophosphamide-treated mice. PT-100 ( $2 \mu\text{g/b.i.d.}$ ), or saline were administered by gavage for the indicated duration. The average absolute neutrophil count for mice not treated with cyclophosphamide was  $194 \times 10^4$  cells/ml as shown by the dotted line.

Figure 8 is a graph depicting the colony formation ability of cells in response to PT-100 in a long-term culture (LTC) assay. Human bone marrow cells were incubated in LTC for 4 weeks in the absence or presence of indicated amounts of PT-100, followed by a 2 week culture in semi-solid medium.

Figure 9 is a graph showing that PT-100 stimulates hematopoiesis in the spleen of normal mice.

- 8 -

Figure 10 is a graph showing that PT-100 stimulates production of G-CSF by human stromal cells.

### **Detailed Description of the Invention**

The invention involves the stimulation of proliferation, differentiation and mobilization  
5 of hematopoietic cells. The invention is useful whenever it is desirable to stimulate the proliferation or differentiation, of or to mobilize, hematopoietic cells. Mobilization of hematopoietic cells is characterized by the enrichment of early progenitor cells in the bone marrow and the recruitment of these cells to the periphery in response to a mobilization agent (e.g. G-CSF, GM-CSF, etc.). The agents useful according to the invention can be used to inhibit  
10 hematopoietic cell deficiencies or to restore hematopoietic and mature blood cell count in subjects with such deficiencies. Such agents also may be used in connection with hematopoietic cell transplants, such as bone marrow or peripheral blood transplants, when used to replenish or create an immune system in a subject. The agents further can be used as an immune booster. The agents also are useful *in vitro* in connection with the culturing of cells for therapeutic and research  
15 uses.

As used herein, subject means humans, nonhuman primates, dogs, cats, sheep, goats, horses, cows, pigs and rodents.

One important aspect of the invention involves restoring or preventing a deficiency in hematopoietic cell number in a subject. Such deficiencies can arise, for example, from genetic  
20 abnormalities, from disease, from stress, from chemotherapy (e.g. cytotoxic drug treatment, steroid drug treatment, immunosuppressive drug treatment, etc.) and from radiation treatment.

The invention is useful in general to restore deficiencies created by hematopoietic cell inhibitors. A hematopoietic cell inhibitor is an exogenously-applied agent (such as a drug or radiation treatment) which causes a decrease in the subject of hematopoietic cells and/or mature  
25 blood cells.

Hematopoietic cells as used herein refer to granulocytes (e.g. promyelocytes, neutrophils, eosinophils and basophils), erythrocytes, reticulocytes, thrombocytes (e.g. megakaryoblasts, platelet-producing megakaryocytes and platelets), lymphocytes, monocytes, dendritic cells and macrophages. Mature blood cells consist of mature lymphocytes, platelets, erythrocytes,  
30 reticulocytes, granulocytes and macrophages. In certain important aspects of the invention, the agents useful according to the invention increase the number of neutrophils, erythrocytes and platelets. In connection with neutrophils, the agents may be used to treat, *inter alia*, drug or

radiation-induced neutropenia, chronic idiopathic neutropenia and cyclic neutropenia.

One important aspect of the invention is restoring in a subject "normal" or "protective" hematopoietic cell levels. A "normal" level as used herein can be a level in a control population, which preferably includes subjects having similar characteristics as the treated individual, such as age. The "normal" level can also be a range, for example, where a population is used to obtain a baseline range for a particular group into which the subject falls. The population can also be divided into groups, such as into quadrants, with the lowest quadrant being individuals with the lowest levels of hematopoietic cells and the highest quadrant being individuals having the highest levels of hematopoietic cells. Thus, the "normal" value can depend upon a particular population selected. Preferably, the normal levels are those of apparently healthy subjects which have no prior history of hematopoietic cell disorders. Such "normal" levels, then can be established as preselected values, taking into account the category in which an individual falls. Appropriate ranges and categories can be selected with no more than routine experimentation by those of ordinary skill in the art. Either the mean or another preselected number within the range can be established as the normal preselected value. Likewise, the level in a subject prior to treatment with a hematopoietic cell inhibitor can be used as the predetermined value.

In general, the normal range for neutrophils is about 1800-7250 per  $\mu\text{l}$  (mean -3650); for basophils 0-150 per  $\mu\text{l}$  (mean -30); for eosinophils 0-700 per  $\mu\text{l}$  (mean -150); for macrophages and monocytes 200-950 per  $\mu\text{l}$  (mean -430); for lymphocytes 1500-4000 per  $\mu\text{l}$  (mean -2500); for erythrocytes  $4.2 \times 10^6$  -  $6.1 \times 10^6$  per  $\mu\text{l}$ ; and for platelets  $133 \times 10^3$  -  $333 \times 10^3$  per  $\mu\text{l}$ . The foregoing ranges are at the 95% confidence level.

In connection with certain conditions, the medical community has established certain preselected values. For example, mild neutropenia is characterized as having a count of between 1000 and 2000 per  $\mu\text{l}$ , moderate neutropenia at between 500 and 1000 per  $\mu\text{l}$  and severe neutropenia at below 500 per  $\mu\text{l}$ . Likewise, in adults, a lymphocyte count at less than 1500 is considered a medically undesirable condition. In children the value is less than 3000. Other preselected values will be readily known to those of ordinary skill in the art. The agents useful according to the invention can be used to establish or to re-establish such preselected values, including normal levels.

Protective levels of hematopoietic cells is the number of cells required to confer clinical benefit to the patient. The required levels can be equal to or less than the "normal levels". Such levels are well known to those of ordinary skill in the art. For example, a protective level of

- 10 -

neutrophils is above 1000, preferably, at least 1500.

According to another aspect of the invention, the agents useful herein can be applied at doses below those which were described in the prior art. In particular, it has been discovered unexpectedly that the agents of the invention can be administered in doses less than 1 mg/kg body weight per day. In particular, the agents of the invention have been used successfully at levels of 0.1 mg/kg body weight per day, which is 10 orders of magnitude below the teachings of the prior art. As will be readily recognized by those of ordinary skill in the art, this has advantages in that less material is required for treatment, thereby lessening any risk of side effects. Likewise, this has advantages in connection with the cost of manufacture of the drug products of the invention.

According to another aspect of the invention, better therapeutic results can be achieved when the agents are applied in multiple doses per day. This finding is unexpected and, additionally, it has been found that there is no added medically useful effect when the agents useful according to the invention are administered for lengthy periods of time. Thus, it has been discovered, unexpectedly, that only very brief periods of treatment are needed to achieve established therapeutic goals.

As described in the examples below, subjects treated with the agents useful according to the invention in 2 doses per day versus 1 dose per day achieved recovery of hematopoietic cells almost 33% faster than subjects receiving only 1 dose per day. Surprisingly, this result did not depend upon the absolute amount of drug given to the subject, but instead related to the number of times the subject was administered the drug. In other words, as shown below, giving twice as much drug, but only once a day, did not speed the recovery of hematopoietic cell number. Thus, an aspect of the invention involves giving the agents useful according to the invention in 2 or 3 doses in an 18 hour period. As used herein, an 18 hour period refers in general to the time during which a subject is awake in any 24 hour period; it is intended to indicate 2 doses per day, 3 doses per day, and the like.

According to still another aspect of the invention, it has been discovered unexpectedly that the agents useful according to the invention need be administered for fewer days than expected according to the prior art. In particular, in the mouse models employed, there was very little difference in the speed of recovery of hematopoietic cell count and in the ability to reestablish normal levels of hematopoietic cells when treatment was 3 days, versus 4 days, versus 5 days. It is believed, therefore, that when applied to humans, a complete drug treatment will involve 7

- 11 -

days or less, more preferably 6 days or less, more preferably 5 days or less, more preferably 4 days or less, and even more preferably 3 days or less. As a result, the invention therefore provides kits which contain complete treatment packages for restoring hematopoietic cell count, which kits are described in greater detail below.

5           According to another aspect of the invention, the time that a subject has an abnormally low level of hematopoietic cells resulting from treatment with a hematopoietic cell inhibitor is shortened. It has been discovered, unexpectedly, that the agents used according to the invention stimulate growth factor production by stromal cells. For example, granulocyte colony stimulating factor (GCSF) production by stromal cells is stimulated. GCSF acts to drive specifically  
10   neutrophil-lineage differentiation. It does not affect the differentiation or proliferation of other committed hematopoietic cells, including other granulocytes, such as eosinophils, basophils, mast cells and macrophages. (It is known to act synergistically, however, *in vitro* with other cytokines to affect proliferation of pluripotent stem cells, though the *in vivo* importance of this observation is not known). Because stromal cells are not rapidly dividing cells and are not generally  
15   adversely impacted by hematopoietic cell inhibitors, the agents useful according to the invention can be applied to subjects substantially simultaneously with or even prior to treatment with a hematopoietic cell inhibitor in order to stimulate stromal cells to produce growth factor which will be readily abundant and helpful in regenerating the hematopoietic cells after treatment by the hematopoietic cell inhibitor. In the prior art, such treatment has been delayed until substantially  
20   after treatment with the hematopoietic cell inhibitor. Substantially simultaneously with, as used herein, means within 24 hours of treatment with the hematopoietic cell inhibitor. Preferably, the agents useful according to the invention are administered within 2 hours of treatment with the hematopoietic cell inhibitor, if they are administered after treatment with the hematopoietic cell inhibitor. If they are administered before treatment with the hematopoietic cell inhibitor, then  
25   they are administered close enough in time to the treatment with the inhibitor so that stromal cell production of growth factor is enhanced in the days immediately following treatment with the hematopoietic cell inhibitor.

          Another aspect of the invention involves treatment of a subject to prepare a subject for subsequent treatment with other agents. It has been discovered, unexpectedly, that the agents  
30   useful according to the invention stimulate the proliferation of primitive, noncommitted hematopoietic progenitor cells, but not directly the differentiation of committed progenitor cells. It is known in the art that such cells may or may not include CD34<sup>+</sup> cells. CD34<sup>+</sup> cells are

- 12 -

immature cells present in blood products, express the CD34 cell surface marker, and are believed to include a subpopulation of cells with the capacity to self-renew and to differentiate into all of the mature blood cell types. Because the agents useful according to the invention stimulate the proliferation of such self-renewing cells, the invention is useful to prepare a subject for treatment  
5 with other exogenous growth factors and cytokines which in turn result in the differentiation of such uncommitted progenitor cells into committed progenitor cells. Likewise, the agents useful according to the invention can be administered to a subject to expand in the subject hematopoietic cells and to mobilize such cells, prior to extracting the cells from the subject for transplantation or re-infusion. Such cells may be used for research purposes or can be treated *ex vivo* or  
10 reintroduced into the subject with or without expansion *in vitro*.

The agents useful according to the invention can be administered in conjunction with exogenous growth factors and cytokines which are specifically selected to achieve a particular outcome. For example, if it is desired to stimulate a particular hematopoietic cell type, then growth factors and cytokines which stimulate proliferation and differentiation of such cell type  
15 are used. Thus, it is known that interleukins-1, 2, 3, 4, 5, 6, 7, 9, 10, 11, 12, 13 and 17 are involved in lymphocyte differentiation. Interleukins 3 and 4 are involved in mast cell differentiation. Granulocyte macrophage colony stimulating factor (GMCSF), interleukin-3 and interleukin-5 are involved in the eosinophil differentiation. GMCSF, macrophage colony stimulating factor (MCSF) and IL-3 are involved in macrophage differentiation. GMCSF, GCSF  
20 and IL-3 are involved in neutrophil differentiation. GMSCF, IL-3, IL-6, IL-11 and TPO are involved in platelet differentiation. Flt3 Ligand is involved in dendritic cell growth. GMCSF, IL-3, and erythropoietin are involved in erythrocyte differentiation. Finally, the self-renewal of primitive, pluripotent progenitor cells capable of sustaining hematopoiesis requires SCF, Flt3 Ligand, G-CSF, IL-3, IL-6 and IL-11. Various combinations for achieving a desired result will  
25 be apparent to those of ordinary skill in the art. Because the agents useful according to the invention stimulate primitive, non-committed hematopoietic progenitor cells, they can be used in connection with any of the foregoing categories of agents to stimulate specifically the proliferation of a particular hematopoietic cell type. The foregoing factors are well known to those of ordinary skill in the art, and most are commercially available.

30 The invention also lends itself to a variety of *in vitro* uses. Hematopoietic progenitor cells are preserved or expanded, or their colony forming unit potential increased, *in vitro*. One benefit that can be obtained according to the invention is the stimulation of hematopoietic progenitor

- 13 -

cells by the agents useful according to the invention. Another benefit that can be obtained is the effect that the agent can have on stromal cells used in *in vitro* culturing of hematopoietic progenitor cells. *In vitro* culturing of hematopoietic cells is often carried out in the presence of stromal cells. Hematopoietic progenitor cells typically will not survive, proliferate or  
5 differentiate for very long periods of time *in vitro* without appropriate growth factor support.

Stromal cell layers are used to supply such growth agents to cultured hematopoietic cells, either by culturing the hematopoietic progenitor cells *in vitro* with such stromal cells or by supplying the hematopoietic progenitor cells with stromal cell-conditioned medium. The agents useful according to the present invention can be used to treat such stromal cells to cause the  
10 stromal cells to manufacture and release growth factors. The incubation of stromal cells with the agents useful according to the invention and in medium is for a period of time sufficient to allow the stromal cells to secrete factors into the medium. The medium then can be used to supplement the culture of hematopoietic progenitor cells and other hematopoietic cells.

The culture of hematopoietic cells is with media which is conventional for culturing cells.  
15 Examples include RPMI, DM, ISCOVES, etc. The conditions for such culturing also are known to those of ordinary skill in the art. The conditions typically refer to a combination of parameters (e.g. temperature, CO<sub>2</sub> and O<sub>2</sub> content, nutritive media, etc.). The time sufficient to increase the number of cells is a time that can be easily determined by a person skilled in the art, and can vary depending on the original number of cells seeded and the amount added of growth factors and  
20 agents useful according to the invention.

The colony forming potential of hematopoietic uncommitted progenitor cells can be increased by *in vitro* culturing of hematopoietic cells. The cells can be obtained from any blood product or organ containing cells of hematopoietic origin. Crude or unfractionated blood products can be enriched for cells having hematopoietic progenitor cell characteristics in ways  
25 well known to those of ordinary skill in the art, prior to or after culture with the agents useful according to the invention.

A particularly important aspect of the invention is in the use of the agents for treatment of neutropenia. A combination of unexpected results makes the invention particularly useful in the treatment of neutropenia. Firstly, the agents according to the invention can stimulate the  
30 proliferation of uncommitted progenitor cells. Secondly, the agents according to the invention also stimulate stromal cells to make GCSF, which is the growth factor critical in the differentiation and production of neutrophils *per se*. Thus, the patient has the dual benefit of

- 14 -

stimulation of progenitor cells and differentiation of those cells into neutrophils using the agents useful according to the invention. Similar effects are shown with erythrocytes and platelets. Thus, treatment to restore neutrophils, erythrocytes and platelets form an independent and distinct aspect of the invention, based on the unexpected findings described above.

5       The invention also involves kits for housing an entire medicinal course of treatment for a hematopoietic cell deficiency such as neutropenia. As discussed above, it has been discovered surprisingly that the number of doses per day and the number of doses overall affect favorably the recovery of hematopoietic cells after treatment with a hematopoietic cell inhibitor. These unexpected findings lend themselves to the development of a medicinal dispenser which houses  
10 an entire medical course of treatment using the agents useful according to the invention. Patient compliance therefore will be enhanced, and an entire prescription can be contained in a single package. Ordinarily, a pharmacist individually fills a dispenser unit with a medicament once the pharmacist receives a doctor's prescription. Because the dispenser of the invention includes an entire medicinal course of treatment and can always include a specific number of solid oral  
15 dosage forms, the package can be pre-filled with the appropriate number of units of medicament for treatment for a particular medical purpose.

The medicinal dispenser is a package defining a plurality of medicinal storage compartments, each compartment for housing an individual unit of medicament. An entire medicinal course of treatment is housed in a plurality of medicinal storage compartments.

20       A package defining a plurality of medicinal storage compartments may be any type of disposable pharmaceutical package or card which holds medicaments in individual compartments. Preferably the package is a blister package constructed from a card, which may be made from stiff paper material, a blister sheet and backing sheet. Such cards are well known to those of ordinary skill in the art.

25       Figure 1 shows a medicinal dispenser (1) for housing a preferred entire medicinal course of treatment for neutropenia. The day indicia (2) indicate which day the individual units of medicament are to be taken. These are marked along a first side of the medicinal package. The dose indicia (3) is marked along a second side of the medicinal package perpendicular to the first side of the medicinal package and indicates the time which the individual unit of medicament  
30 should be taken. The unit doses (4) are contained in the dispenser which is a blister pack. This particular package shows a 5 day course of treatment, with 2 doses per day.

The pharmaceutical preparations, as described above, are administered in effective



- 15 -

amounts. The effective amount will depend upon the mode of administration, the particular condition being treated and the desired outcome. It will also depend upon, as discussed above, the stage of the condition, the age and physical condition of the subject, the nature of concurrent therapy, if any, and like factors well known to the medical practitioner. For therapeutic applications, it is that amount sufficient to achieve a medically desirable result. In some cases this is any increase in hematopoietic cell count or mature blood cell count. In other cases, it will be an increase to a preselected level.

The invention is useful in one aspect to ameliorate the effects of treatment with a hematopoietic cell inhibitor. If the agents are used prophylactically, they can decrease the amount of hematopoietic cells that would be lost in the subject versus the amount lost if the subject were treated with the inhibitor but not with the agent. If used prophylactically or acutely, the agents can shorten the time for recovery of a hematopoietic cell-type to at least protective levels, and preferably to normal levels, versus the length of time which would pass before protective or normal levels were achieved if the subject were treated with the inhibitor but not with the agent.

Generally, doses of active compounds of the present invention would be from about 0.01 mg/kg per day to less than 1 mg/kg per day. A variety of administration routes are available. The methods of the invention, generally speaking, may be practiced using any mode of administration that is medically acceptable, meaning any mode that produces effective levels of the active compounds without causing clinically unacceptable adverse effects. Such modes of administration include oral, rectal, topical, nasal, interdermal, or parenteral routes. The term "parenteral" includes subcutaneous, intravenous, intramuscular, or infusion. Intravenous or intramuscular routes are not particularly suitable for long-term therapy and prophylaxis. They could, however, be preferred in emergency situations. Oral administration is preferred for the convenience to the patient as well as the dosing schedule. See Remington's Pharmaceutical Sciences, 18th edition, 1990, pp 1694-1712; incorporated by reference). Those of skill in the art can readily determine the various parameters and conditions for producing dosages without resort to undue experimentation.

Compositions suitable for oral administration may be presented as discrete units, such as capsules, tablets, lozenges, each containing a predetermined amount of the active agent. Other compositions include suspensions in aqueous liquids or non-aqueous liquids such as a syrup, elixir or an emulsion.

Preparations for parenteral administration include sterile aqueous or non-aqueous

- 16 -

solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, 5 Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like. Lower doses will result from other forms of administration, such as intravenous administration. In the 10 event that a response in a subject is insufficient at the initial doses applied, higher doses (or effectively higher doses by a different, more localized delivery route) may be employed to the extent that patient tolerance permits. Multiple doses per day are contemplated.

The agents may be combined, optionally, with a pharmaceutically-acceptable carrier. The term "pharmaceutically-acceptable carrier" as used herein means one or more compatible solid 15 or liquid filler, diluents or encapsulating substances which are suitable for administration into a human. The term "carrier" denotes an organic or inorganic ingredient, natural or synthetic, with which the active ingredient is combined to facilitate the application. The components of the pharmaceutical compositions also are capable of being co-mingled with the molecules of the present invention, and with each other, in a manner such that there is no interaction which would 20 substantially impair the desired pharmaceutical efficacy.

When administered, the pharmaceutical preparations of the invention are applied in pharmaceutically-acceptable amounts and in pharmaceutically-acceptably compositions. Such preparations may routinely contain salt, buffering agents, preservatives, compatible carriers, and optionally other therapeutic agents. When used in medicine, the salts should be pharmaceutically 25 acceptable, but non-pharmaceutically acceptable salts may conveniently be used to prepare pharmaceutically-acceptable salts thereof and are not excluded from the scope of the invention. Such pharmacologically and pharmaceutically-acceptable salts include, but are not limited to, those prepared from the following acids: hydrochloric, hydrobromic, sulfuric, nitric, phosphoric, maleic, acetic, salicylic, citric, formic, malonic, succinic, and the like. Also, pharmaceutically- 30 acceptable salts can be prepared as alkaline metal or alkaline earth salts, such as sodium, potassium or calcium salts.

Other delivery systems can include time-release, delayed release or sustained release

- 17 -

delivery systems. Such systems can avoid repeated administrations of the agent, increasing convenience to the subject and the physician. Many types of release delivery systems are available and known to those of ordinary skill in the art. They include polymer base systems such as poly(lactide-glycolide), copolyoxalates, polycaprolactones, polyesteramides, polyorthoesters, polyhydroxybutyric acid, and polyanhydrides. Microcapsules of the foregoing polymers containing drugs are described in, for example, U.S. Patent 5,075,109. Delivery systems also include non-polymer systems that are: lipids including sterols such as cholesterol, cholesterol esters and fatty acids or neutral fats such as mono- di- and tri-glycerides; hydrogel release systems; sylastic systems; peptide based systems; wax coatings; compressed tablets using conventional binders and excipients; partially fused implants; and the like. Specific examples include, but are not limited to: (a) erosional systems in which the agent is contained in a form within a matrix such as those described in U.S. Patent Nos. 4,452,775, 4,667,014, 4,748,034 and 5,239,660 and (b) difusional systems in which an active component permeates at a controlled rate from a polymer such as described in U.S. Patent Nos. 3,832,253, and 3,854,480. In addition, pump-based hardware delivery systems can be used, some of which are adapted for implantation.

Use of a long-term sustained release implant may be particularly suitable for treatment of chronic conditions. Long-term release, as used herein, means that the implant is constructed and arranged to deliver therapeutic levels of the active ingredient for at least 30 days, and preferably 60 days. Long-term sustained release implants are well-known to those of ordinary skill in the art and include some of the release systems described above.

#### Examples

We have demonstrated in a series of *in vivo* studies that the agent ValboroPro (PT-100), has the ability to shorten myelosuppression caused by chemotherapy in mice. In these studies, mice were injected intraperitoneally with a sublethal dose of 220 mg/kg cyclophosphamide (Day 1). This treatment reproducibly induced a nadir in blood cell counts by Day 4. After 72 hours (Day 3) mice were divided into 3 groups. One group received PT-100, at the concentrations indicated, by gavage or by subcutaneous administration (s.c.), one group received G-CSF by s.c. injections and the third group received saline as a control, either by oral gavage or by s.c. injections. G-CSF was used at 0.04 ug/dose (4µg/kg/day) which is the dose frequently used in published reports studying the G-CSF effects in mice and is also the equivalent dose used in cancer patients. All administrations were performed twice daily (b.i.d.) for 5 consecutive days or as indicated. Blood samples were taken from individual mice on Day 4-8, and in some

- 18 -

experiments on Days 13 or 17. At each time point four or five test animals were sampled. Total and differential white blood cell counts of Gimsa-stained blood smears were performed.

#### **PT-100 Dose Response For Regeneration of Neutrophil**

For data presented in Figure 2, cyclophosphamide treated mice received 0.1  $\mu$ g, 2  $\mu$ g or 5  $\mu$ g/b.i.d. of PT-100 or saline by oral gavage twice daily for 5 consecutive days starting on Day 3 post cyclophosphamide treatment and continuing through Day 7. In mice that received 2 or 5  $\mu$ g/b.i.d. PT-100 recovery of neutrophils reproducibly preceded recovery of saline treated mice by 1 or 2 days, while 0.1  $\mu$ g /b.i.d. of PT-100 did not significantly enhance neutrophil recovery over saline. Normal levels of absolute neutrophil counts (ANC) were reached on Day 5 for mice receiving 2  $\mu$ g or 5  $\mu$ g/b.i.d. of PT-100, while saline treated mice did not reach normal levels until Day 7. On Day 5 mice had received a total of 4 doses of PT-100 (on Days 3 and 4). Additional administration of PT-100 on Days 5, 6 and 7 caused a further increase in ANC.

The effect of PT-100 on neutrophil recovery when administered by s.c. route was very similar to that seen when administered orally. For data shown in Figure 3 mice were injected s.c. with doses of PT-100 ranging from 1 to 20  $\mu$ g/b.i.d. for 5 days and blood cell counts determined on Days 4 through 8, and on Day 17. For mice receiving 5  $\mu$ g, 10  $\mu$ g, or 20  $\mu$ g/b.i.d. PT-100, neutrophil recovery was accelerated over that observed in the saline treated mice. A dose of 1  $\mu$ g/b.i.d. PT-100 did not show much effect. After termination of treatment with PT-100, in conclusion, PT-100 accelerates neutrophil regeneration in cyclophosphamide treated mice.

#### **Comparison Of PT-100 And G-CSF Effects On Neutrophil Regeneration**

G-CSF is currently used to accelerate neutrophil recovery in cancer patients undergoing chemotherapy. The effects of G-CSF in mice are well established and can be used as a reference for elucidating the mechanism by which PT-100 stimulates hematopoiesis in mice. Figure 4 shows data from an experiment in which the effects of PT-100 and G-CSF on neutrophil regeneration are compared. Cyclophosphamide treated mice were administered 2  $\mu$ g/b.i.d. of PT-100 by gavage or 0.04  $\mu$ g/b.i.d. of G-CSF (the dose equivalent used in patients and most commonly used in published reports for murine studies) by subcutaneous injections for 5 consecutive days starting on Day 3. Blood cell counts were performed on Days 4-8, and on Day 13.

PT-100 and G-CSF treated mice stimulated neutrophil regeneration to a similar level during the treatment period. After treatment was stopped, ANC decreased to normal counts by Day 13. Although PT-100 has a very similar effect on neutrophil reconstitution, the mechanism

- 19 -

of action is different from that of G-CSF. Not only does PT-100 target a different cellular receptor (CD 26), it also has been shown to stimulate growth of early human hematopoietic progenitor cells which are not affected by G-CSF.

#### **Dose Numbering Of PT-100 Administration**

5 To determine the dose numbering of administration for optimal recovery of neutrophils, PT-100, at indicated doses, was administered s.c. to cyclophosphamide treated mice, either once or twice per day over a five day period, starting on Day 3 post cyclophosphamide treatment. As shown in Figure 5, for both doses, a twice daily administration resulted in a faster rate of neutrophil recovery to higher neutrophil levels than once per day administration.

#### **Duration of PT-100 Administration**

10 In the experiments described above mice had been treated with PT-100 for 5 consecutive days. To determine whether a shorter period of treatment with PT-100 was sufficient for the recovery of neutrophils 5  $\mu$ g, 2  $\mu$ g, or 1  $\mu$ g/b.i.d. (six hours apart) of PT-100 was administered to cyclophosphamide treated mice by gavage for 1,2,3, or 5 days starting on Day 3 post  
15 cyclophosphamide treatment. Blood counts were obtained on days 4 through 8.

Administration of PT-100 for one day was sufficient to cause an accelerated reconstitution of neutrophils over saline treated animals. However, additional administrations of PT-100 for 2 or 3 days increased the rate of recovery even further. Data for the 5  $\mu$ g dose are shown in Figure 6.

20 Continued administration of PT-100 for a total of 4 or 5 days does not significantly increase the rate of neutrophil recovery or the ANC over that achieved with 3 day administrations (data for 2  $\mu$ g/b.i.d. are shown in Figure 7).

Results shown in Figures 6 and 7 indicate that the PT-100 effect on the regeneration of neutrophils occurs early during treatment and continues until ANC between 1000 and 1400 are  
25 achieved. Repeated administrations affect the kinetics of neutrophil restoration during the early period but does not significantly alter the ANC reached after 3 days of administration.

In conclusion, PT-100 accelerates neutrophil reconstitution over that seen with saline even after a one day of treatment. An accelerated reconstitution of neutrophils is obtained with each additional day of treatment for up to three days. A fourth or fifth day of treatment did not  
30 significantly increase ANC or the kinetics of reconstitution.

#### **Human Hematopoietic Cell Responses *In Vitro***

Hematopoiesis is sustained by a pool of hematopoietic stem cells (HSCs) that can self-

- 20 -

renew and differentiate into hematopoietic progenitor cells (HPCs). HPCs are committed to specific lineages which can be identified based on their colony morphology when grown in semi-solid media *in vitro*, typically over a 2 week period. The colonies grown in the semi-solid colony assay are functionally defined as colony- or burst- forming units and include BFU-E and CFU-E (cells committed to the erythroid lineage), CFU-GM (cells committed to the granulocytic/monocytic lineage), BFU-MK and CFU-MK (cells committed to the megakaryocyte lineage) and CFU-GEMM (multipotent progenitors). Although the semi-solid colony assay is a valuable tool to identify factors, such as G-CSF, which affect terminal differentiation, it does not assess the proliferative potential or self renewing properties of the primitive hematopoietic progenitor cells (PHPCs) (Dexter, T.A. et al., *Acta Hemat.* 62:299-305 (1979); Chen, B.P. et al., *Immunological Reviews*:157:41-51 (1997)).

An assay to evaluate the effect of a compound or of growth factors on PHPCs was first described by Dexter (Dexter T.M. et al, *J. Cell. Physiol.* 91:335-344 (1977)), and combines the Long-Term Culture (LTC) with the semi-solid colony assay. LTC is initiated over a pre-formed stromal cell layer which provides the necessary hematopoietic growth factors. It has been used extensively for the *in vitro* examination of murine and human hematopoiesis and to evaluate the ability of test compounds to generate LTC-ICs.

The effect of PT-100 on growth of human hematopoietic cells was examined in the 2 week CFU and the 4 and 5 week LTC assays using human bone marrow, apheresed peripheral blood or umbilical cord blood cells. PT-100 did not stimulate the generation of CFUs in the 2 week semi-solid assay, indicating that PT-100 does not affect the differentiation of committed progenitor cells into mature blood cells. It also suggests, that the mechanism and the cellular targets for PT-100 for the stimulation of neutrophil regeneration *in vivo* is different from that of G-CSF which has been shown to stimulate CFU formation in this assay. In the LTC assays, which test for effects on early progenitor cells, PT-100 significantly increased the growth of very early progenitor cells from all three cell sources. Moreover, the data suggest that the effect of PT-100 is on PHPCs as increases in LTC-ICs were observed at 4 weeks (Figure 8) 5 weeks and 6 weeks (data not shown) in culture. At this time less primitive hematopoietic progenitor cells have undergone terminal differentiation and lost the ability to form colonies in semi-solid cultures.

For the LTC assays, CD34<sup>+</sup> cells were isolated by positive selection from human bone marrow cells, apheresed peripheral blood or umbilical cord blood using a MAC separation system. To establish a stromal feeder layer, human bone marrow cells were cultured in Myelocult

- 21 -

long term culture medium for 2 weeks. One day prior to use, the adherent stromal cells were cultured overnight with indicated concentrations of PT-100 in LTC medium and irradiated. Isolated CD34<sup>+</sup> cells were overlaid onto the stromal cell layer and incubated for 30 days in the absence or presence of indicated amounts of PT-100. Medium and PT-100 was exchanged every  
5 three days thereafter. At the end of the culture period the culture was assayed for progenitor cells by plating in semi-solid medium (methylcellulose) supplemented with growth factors (Stem Cell Factor, GM-CSF, IL-3 and Erythropoietin).

The total number of myeloid, erythroid, blast forming and multilineage clonogenic progenitors (colonies CFU-GM, CFU-E, BFU-E and CFU-GEMM, respectively) were determined  
10 after 14 days in methylcellulose culture.

Data showing in Figure 8 for a human bone marrow culture indicate that during a 4 week LTC assay, PT-100 increased, in a dose dependent manner, the number of clonogenic progenitors which are able to form colonies in semi-solid medium. This suggests that PT-100 stimulates growth of primitive hematopoietic progenitor cells.

15 In similar fashion CD34<sup>+</sup> cells purified from apheresed peripheral blood or umbilical cord blood were cultured on irradiated primary stromal cells for 30 days. As had been observed with bone marrow cells, PT-100 increased the number of 4 and 5 week LTC-ICs from peripheral and umbilical cord blood to very similar levels, indicating the PT-100 is able to stimulate primitive hematopoietic progenitor cell growth from these cell sources as well (data not shown).

20 **PT-100 Does Not Stimulate Differentiation of Committed Progenitor Cells**

Human bone marrow cells were enriched for CD34<sup>+</sup> cells and 200 CD34<sup>+</sup> cells per well were incubated in serum free *x-vivo* 15 medium (Biowhittaker) with or without the indicated concentrations of PT-100 for 4 hours at 37°C. The pre-incubated CD34<sup>+</sup> cells were added to 0.9% methylcellulose in Iscove's MDM containing sub-optimal concentrations of recombinant  
25 human growth factors (5 ng/ml Stem Cell Factor, 1 ng/ml GM-CSF, 1 ng/ml IL-3, 0.3 units/ml Erythropoietin (Stem Cell Technologies Vancouver, BC). PT-100 was added to the medium at the same concentrations used for the pre-incubation. The methylcellulose mixture was plated in duplicate in 35 mm dishes and incubated for 14 days at 37°C. Progenitor colonies (CFU-E, CFU-GM, CFU-GEMM and BFU-E) were counted under an inverted microscope. PT-100 did not  
30 stimulate differentiation of these committed progenitor cells.

**Stimulation of Hematopoiesis in the Spleen of Normal Mice**

6-8 week old female BALB/c mice were administered either saline or PT-100 twice daily

- 22 -

for 5 days at the indicated doses via either subcutaneous injection or oral gavage. On the sixth day the animals were sacrificed and their spleens were excised using sterile procedures. The spleens were disrupted to produce single cell suspensions which were subsequently treated with a solution of Tris ammonium chloride (pH 7.2) to lyse erythrocytes. The resulting splenocyte populations were in a hemocytometer and resuspended at  $5 \times 10^6$  cells/mL in Iscove's Modified Eagles medium (IMDM) supplemented with 2% heat inactivated fetal calf serum. 0.3 mL of each splenocyte solution was added to 3 mL of Methocult™MGF M3434 (Stem Cell Technologies, Vancouver, BC, Canada), a methylcellulose medium containing recombinant cytokines used for colony assays of murine progenitor cells. The medium was vigorously mixed and then 1.1 mL of the mixture was placed in duplicate onto sterile 35 mm diameter culture dishes, resulting in  $5 \times 10^5$  splenocytes/plate. The plated cells were incubated at 37°C under humidified conditions in 95% air/5% CO<sub>2</sub> for 7 days. CFU-E were enumerated as per the manufacturers specifications after 2 days, while BFU-E, CFU-GM and CFU-GEMM were enumerated after 7 days. For each mouse, the absolute CFU/spleen were calculated using the total splenocyte count determined in the hemocytometer. The data shown in Figure 9 represents the mean  $\pm$  SD CFU/spleen from 3 mice in each dosing group. PT-100 stimulated hematopoiesis for all progenitor colony types tested.

#### **PT-100 Induces Production of G-CSF From Human Bone Marrow Stromal Cells**

Mononuclear cells were purified from bone marrow and cultured long-term culture medium, (Stem Cell Technologies, Inc., Vancouver, B.C.) for 2 weeks, with a single feeding of fresh medium after 1 week. The established stromal cells were removed by trypsin-EDTA digest and seeded into a 35 mm tissue culture plate at  $10^6$  cells per well in 1 ml of medium containing  $10^{-5}$  M PT-100 or medium alone as control. Culture media were collected on day 1. Supernatants were assayed for human G-CSF using a Quantikine high sensitivity immuno-assay kit (R + D Systems, Minneapolis, MN). Figure 10 depicts the effect of PT-100 on the production of G-CSF by cultured human stromal cells. PT-100 stimulates production of G-CSF by such cells.

The manufacture of L-VAL-R-boroPro is described in a number of published procedures (Kelly, T.A., et al. *J. Am. Chem. Soc.* 1993. 115:12537-12638; Coutts, S.J., et al., *J. Med. Chem.* 1996. 39:2087-2094; Beak, P., et al., *Tetrahedron Letters*, 1989, 30:1197; Bean, F.R., et al., *J. Amer. Chem. Soc.* 1932. 54:4415). Pure isomers are preferred. See also US Patents 4,935,493 and 5,462,928, the disclosures of which are incorporated here by reference.

While the invention has been described with respect to certain embodiments, it should be



- 23 -

appreciated that many modifications and changes may be made by those of ordinary skill in the art without departing from the spirit of the invention. It is intended that such modifications, changes, and equivalents fall within the scope of the following claims.

- 24 -

We claim:

1. A method for treating a subject to stimulate hematopoiesis in the subject, comprising:  
administering to a subject in need of such treatment an amount of an agent effective to  
increase the number of hematopoietic cells or mature blood cells in the subject, wherein the  
5 amount is less than 1 mg/kg body weight per day and wherein the agent is a compound of  
Formula I.
2. The method of claim 1, wherein the subject has an abnormally low level of hematopoietic  
cells and wherein the agent is administered in an amount effective to restore levels of a  
10 hematopoietic cell-type to a preselected normal or protective level.
3. The method of claim 1, wherein the subject is administered at least two doses of the agent  
in an 18 hour period.
- 15 4. The method of claim 2, wherein the subject is neutropenic and wherein said amount is  
effective to restore a preselected normal or protective level of neutrophils.
5. The method of claim 2, wherein the subject has abnormally low levels of erythrocytes and  
wherein said amount is effective to restore a preselected normal or protective level of  
20 erythrocytes.
6. The method of claim 2, wherein the subject has abnormally low levels of platelets and  
wherein said amount is effective to restore a preselected normal or protective level of platelets.
- 25 7. The method of claims 1, 2, 3, 4, 5 or 6, wherein the agent is ValBoroPro.
8. A method for shortening the time that a subject has an abnormally low level of  
hematopoietic or mature blood cells resulting from treatment with a hematopoietic cell inhibitor,  
comprising:  
30 administering to a subject in need of such treatment an agent in an amount effective to  
increase the number of hematopoietic cells or mature blood cells in the subject,  
wherein the administration of the agent is prior to or substantially simultaneous with

- 25 -

administration of the hematopoietic cell inhibitor, and wherein the agent is a compound of Formula I.

9. The method of claim 8, wherein the administration of the hematopoietic cell inhibitor  
5 causes an abnormally low level of hematopoietic cells in the subject and wherein the agent is administered in an amount effective to restore levels of a hematopoietic cell-type to a preselected normal or protective level.

10. The method of claim 9, wherein the subject is administered at least two doses of the agent  
10 in an 18 hour period.

11. The method of claim 9, wherein the hematopoietic cell inhibitor reduces in the subject  
neutrophils and wherein the amount is effective to restore in the subject a preselected normal or  
protective level of neutrophils.

12. The method of claim 9, wherein the hematopoietic cell inhibitor reduces in the subject  
erythrocytes and wherein the amount is effective to restore in the subject a preselected normal or  
protective level of erythrocytes.

13. The method of claim 9, wherein the hematopoietic cell inhibitor reduces in the subject  
platelets and wherein the amount is effective to restore in the subject a preselected normal or  
protective level of platelets.

14. The method of claims 8, 9, 10, 11, 12 or 13, wherein the amount is less than 1 mg/kg body  
25 weight per day.

15. The method of claims 8, 9, 10, 11, 12 or 13, wherein the agent is ValBoroPro.

16. A method for preparing a subject for treatment with a hematopoietic cell inhibitor,  
30 comprising:

administering to the subject prior to the subject receiving the hematopoietic cell inhibitor  
an agent in an amount effective to stimulate in the subject production of growth factor, wherein

- 26 -

the agent is a compound of Formula I.

17. The method of claim 16, wherein the growth factor is granulocyte colony stimulating factor.

5

18. The method of claim 16, wherein the amount is less than 1 mg/kg body weight per day.

19. The method of claim 16, wherein the agent is administered to the subject by administering at least two doses of the agent in an 18 hour period.

10

20. The method of claim 16, wherein the agent stimulates stromal cell production of growth factor.

21. The method of claims 16, 17, 18, 19 or 20, wherein the agent is ValBoroPro.

15

22. A method for treating a subject to increase the number of hematopoietic cells in the subject comprising:

administering to a subject in need of such treatment an amount of an agent effective to increase hematopoietic cells in the subject,

20

wherein the agent is administered in a first regimen consisting of 2 doses or 3 doses in an 18 hour period, and wherein the agent is a compound of Formula I.

25

23. The method of claim 22, wherein the agent is administered in a second regimen consisting of 2 doses or 3 doses in an 18 hours period, and wherein, the second regimen is separate in time from the first regimen.

30

24. The method of claim 23, wherein the agent is administered in a third regimen consisting of 2 doses or 3 doses in an 18 hours period, and wherein, the third regimen is separate in time from the first and second regimens.

25. The method of claim 24, wherein the agent is administered in a fourth regimen consisting of 2 doses or 3 doses in an 18 hours period, and wherein, the fourth regimen is separate in time

- 27 -

from the first, second, and third regimens.

26. The method of claim 25, wherein the agent is administered in a fifth regimen consisting of 2 doses or 3 doses in an 18 hours period, and wherein, the fifth regimen is separate in time  
5 from the first, second, third and fourth regimens.

27. The method of claims 22, 23, 24, 25, or 26, wherein the subject has abnormally low neutrophils and, wherein said amount is effective to restore a preselected normal or protective level of neutrophils.

10

28. The method of claims 22, 23, 24, 25, or 26, wherein each regimen is less than 1 mg/kg body weight per day.

29. The method of claims 22, 23, 24, 25, or 26, wherein the agent is ValBoroPro.

15

30. A method for preparing a subject's cells for reintroduction into the subject comprising administering to the subject an agent in an amount effective to stimulate hematopoietic cells,

then collecting hematopoietic cells from the subject, and  
20 reintroducing said collected cells into the subject,  
wherein the agent is a compound of Formula I.

31. The method of claim 30, further comprising *ex-vivo* culturing the collected cells in the presence of an amount of the agent effective to stimulate proliferation of said collected cells.

25

32. The method of claim 30, wherein the concentration of the agent in medium surrounding the collected cells is less than  $10^{-8}$  moles per liter.

33. The method of claims 30, 31, or 32, wherein the agent is ValBoroPro.

30

34. A method for stimulating growth factor production by stromal cells comprising:  
contacting the stromal cells with an agent in an amount effective to stimulate growth

- 28 -

factor production by the stromal cells, wherein the agent is a compound of Formula I.

35. The method of claim 34, wherein the stromal cells are an *in vitro* layer of stromal cells for supporting cell growth and further comprising culturing said stem cells in the presence of said  
5 stromal cells.

36. The method of claim 34, wherein the stromal cells are *in vivo* in a subject.

37. The method of claim 34, wherein the growth factor is granulocyte colony stimulating  
10 factor.

38. The method of claim 36, wherein the agent is administered to the subject in an amount less than 1 mg/kg per day.

15 39. The method of claims 34, 35, 36, 37 or 38, wherein the agent is ValBoroPro.

40. The method of claim 35, wherein the stem cells are cultured in an environment free of exogenously added growth factors.

20 41. The method of claim 35, wherein the stem cells are cultured in an environment free of exogenously added granulocyte colony stimulating factor

42. A kit for treating a subject to ameliorate the effects resulting from treatment with a hematopoietic cell inhibitor, comprising a package containing:

25 a first dosage of an agent and instructions for treating the subject with the agent, substantially simultaneous with or prior to treatment with the hematopoietic cell inhibitor, and

a second dosage of the agent and instructions for treating the subject with the agent, only after treatment with the hematopoietic cell inhibitor,

wherein said dosages are in effective amounts and,

30 wherein the agent is a compound of Formula I.

43. The kit of claim 42, wherein the second dosage is between 2 and 7 regimens, each of said

- 29 -

regimens consisting of two or three doses of the agent for administration in an 18 hour period.

44. The kit of claim 42, wherein the second dosage is less than 1 mg/kg body weight per day.

5 45. The kit of claims 42, 43, or 44, wherein the agent is ValBoroPro.

46. The kit of claims 42, 43, or 44, wherein the kit is for treating or inhibiting neutropenia.

47. The kit of claim 45, wherein the kit is for treating or inhibiting neutropenia.

10

48. The kit of claims 42, 43 or 44, wherein the kit is for treating or inhibiting a platelet deficiency.

49. The kit of claim 45, wherein the kit is for treating or inhibiting a platelet deficiency.

15

50. The kit of claims 42, 43, or 44, wherein the kit is for treating or inhibiting an erythrocyte deficiency.

51. The kit of claim 45, wherein the kit is for treating or inhibiting an erythrocyte deficiency.

20

52. A kit for treating a subject having an abnormally low level of hematopoietic cells or mature blood cells comprising:

a packing containing a complete dosage for restoring at least protective levels of a hematopoietic cell or a mature blood cell type, the package consisting essentially of

25 a first dosage of an agent in an effective amount for administration to the subject during a first 18 hour period of a first day,

a second dosage of an agent in an effective amount for administration to the subject during a second 18 hour period of a second day,

30 optionally a third dosage of the agent in an effective amount for administration to the subject during a third 18 hour period of a third day,

optionally a fourth dosage of the agent in an effective amount for administration to the subject during a fourth 18 hour period of a fourth day,

- 30 -

optionally a fifth dosage of the agent in an effective amount for administration to the subject during a fifth 18 hour period of a fifth day,

optionally a sixth dosage of the agent in an effective amount for administration to the subject during a sixth 18 hour period of a sixth day,

5 optionally a seventh dosage of the agent in an effective amount for administration to the subject during a sixth 18 hour period of a seventh day,

wherein the agent is a compound of Formula I and wherein the agent is present in an amount effective to restore a preselected normal or protective level of hematopoietic or mature blood cell-type.

10

53. The kit of claim 52, consisting essentially of only the first, second, third, fourth and fifth dosages.

54. The kit of claim 52, consisting essentially of only the first, second, third and fourth  
15 dosages.

55. The kit of claims 52, 53 or 54, wherein each of the dosages consists of 2 or 3 doses of the agent for each respective 18 hour period.

20 56. The kit of claims 52, 53 or 54, wherein the amount of each dosages is less than 1 mg/kg body weight per day.

57. The kit of claims 52, 53, 54, or 55, wherein the agent is ValBoroPro.

25 58. A kit for ameliorating on a subject the effects of a hematopoietic cell inhibitor comprising a package containing:

a first dosage of an agent for administration to a subject during a first day, and

a second dosage of the agent for administration to a subject during a second day,

optionally a third dosage of the agent in an effective amount for administration to the

30 subject during a third 18 hour period of a third day,

optionally a fourth dosage of the agent in an effective amount for administration to the

subject during a fourth 18 hour period of a fourth day,



- 31 -

optionally a fifth dosage of the agent in an effective amount for administration to the subject during a fifth 18 hour period of a fifth day,

optionally a sixth dosage of the agent in an effective amount for administration to the subject during a sixth 18 hour period of a sixth day,

5 optionally a seventh dosage of the agent in an effective amount for administration to the subject during a sixth 18 hour period of a seventh day,

wherein the combined dosages are in amounts effective to ameliorate the effects in the subject of the hematopoietic cell inhibitor and wherein each dosage is less than 1 mg/kg body weight per day.

10

59. The kit of claim 58, consisting essentially of only the first, second, third, fourth and fifth dosages.

60. The kit of claim 58, consisting essentially of only the first, second, third, and fourth  
15 dosages.











61. The kit of claims 58, 59, or 60, wherein each of the dosages consists of two or three doses of the agent for each respective day.

20 62. The kit of claims 58, 59, or 60, wherein the agent is ValboroPro.

63. The kit of claim 61, wherein the agent is ValboroPro.

1/6

1

DAY	Dose AM	Dose PM
1		
2		
3		
4		
5		

2 3 4

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8

5 / 6




FIG. 9

6/6




FIG. 10

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/09652

A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 6 A61K31/69 A61K38/55

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 98 50046 A (TUFTS COLLEGE) 12 November 1998 (1998-11-12)  the whole document	1-4, 7, 22-29, 34, 36, 38, 39, 42-63
P, X	WO 98 50066 A (TUFTS COLLEGE) 12 November 1998 (1998-11-12)  the whole document	22-27, 29, 34, 36, 39, 42-63

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance  
 "E" earlier document but published on or after the international filing date  
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
 "O" document referring to an oral disclosure, use, exhibition or other means  
 "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.  
 "&" document member of the same patent family

Date of the actual completion of the international search

15 October 1999

Date of mailing of the international search report

21/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Hoff, P



## INTERNATIONAL SEARCH REPORT

International Application No

PC./US 99/09652

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94 03055 A (US HEALTH ;UNIV TUFTS (US)) 17 February 1994 (1994-02-17) cited in the application  the whole document ----	8-13, 15, 22-27, 29, 34, 36, 39, 42-63
X	WO 95 11689 A (TUFTS COLLEGE ;BACHOVCHIN WILLIAM W (US)) 4 May 1995 (1995-05-04) the whole document ----	42-63
A		1-41
X	WO 93 08259 A (NEW ENGLAND MEDICAL CENTER INC ;UNIV TUFTS (US)) 29 April 1993 (1993-04-29) the whole document ----	42-63
A		1-41
A	BRISTOL L A ET AL: "Inhibition of CD26 enzyme activity with pro-boropro stimulates rat granulocyte/macrophage colony formation and thymocyte proliferation in vitro" BLOOD, vol. 85, no. 12, 15 June 1995 (1995-06-15), pages 3602-3609, XP002092855 ISSN: 0006-4971 the whole document ----	1-41
A	COUTTS S J ET AL: "Structure-activity relationships of boronic acid inhibitors of dipeptidyl peptidase IV. 1. Variation of the P2 position of Xaa -boroPro dipeptides" JOURNAL OF MEDICINAL CHEMISTRY, vol. 39, no. 39, 10 May 1996 (1996-05-10), pages 2087-2094, XP002092856 ISSN: 0022-2623 the whole document ----	1-41
A	WO 98 00439 A (TUFTS COLLEGE ;BACHOVCHIN WILLIAM W (US)) 8 January 1998 (1998-01-08) the whole document ----	1-63
P, A	WO 99 16864 A (POINT THERAPEUTICS INC) 8 April 1999 (1999-04-08) the whole document -----	1-63

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC/US 99/09652

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9850046 A	12-11-1998	AU 7269198 A	27-11-1998
WO 9850066 A	12-11-1998	AU 7274798 A	27-11-1998
WO 9403055 A	17-02-1994	AU 4794393 A	03-03-1994
WO 9511689 A	04-05-1995	NONE	
WO 9308259 A	29-04-1993	CA 2121369 A	29-04-1993
		EP 0610317 A	17-08-1994
		JP 7504158 T	11-05-1995
		US 5462928 A	31-10-1995
WO 9800439 A	08-01-1998	AU 3792797 A	21-01-1998
		CA 2258038 A	08-01-1998
		EP 0938498 A	01-09-1999
WO 9916864 A	08-04-1999	AU 9588798 A	23-04-1999